

Contact : JOLYON PRIDDING
4RECYCLING LTD
CONTROL HOUSE
A1 BUSINESS PARK
KNOTTINGLEY ROAD
KNOTTINGLEY WF11 0BU
Tel. : 01132 322400

V537

Please quote the above code for all enquiries

Client :

Sample Matrix : Agricultural Soil

Laboratory Reference

Card Number 05659/25

Date Received 14-Apr-25

Date Reported 22-Apr-25

SOIL ANALYSIS REPORT

Laboratory Sample Reference	Field Details			Index			mg/l (Available)		
	No.	Name or O.S. Reference with Cropping Details	Soil pH	P	K	Mg	P	K	Mg
25291/25	1	BLAENYCOED 1 <i>No cropping details given</i>	6.5	3	3	3	42.0	272	127
25292/25	2	BLAENYCOED 2 <i>No cropping details given</i>	5.9	3	3	3	38.0	394	102
25293/25	3	BLAENYCOED 3 <i>No cropping details given</i>	6.0	3	3	2	35.6	270	85
25294/25	4	BLAENYCOED 4 <i>No cropping details given</i>	6.0	4	3	3	54.6	243	114
25295/25	5	BLAENYCOED 7 <i>No cropping details given</i>	5.6	3	2+	3	27.2	212	126
25296/25	6	BLAENYCOED 8 <i>No cropping details given</i>	5.9	2	2-	2	25.2	147	97

If general fertiliser and lime recommendations have been requested, these are given on the following sheets.

The analytical methods used are as described in DEFRA Reference Book 427

The index values are determined from the AHDB Fertiliser Recommendations RB209 9th Edition.

Released by*Katie Dunn*..... On behalf of NRM Date *22/04/25*.....

Independently Analysed by **NRM**, part of the **Cawood Group**, Coopers Bridge, Braziers Lane, Bracknell, Berkshire RG42 6NS

Tel +44 (0) 1344 886338 Fax: +44 (0) 1344 890972 Email: enquiries@nrm.uk.com www.nrm.uk.com

DATE 22nd April 2025
 SAMPLES FROM

SAMPLED BY

Report reference 05659/25

JOLYON PRIDDING
 4RECYCLING LTD
 CONTROL HOUSE
 A1 BUSINESS PARK
 KNOTTINGLEY ROAD
 KNOTTINGLEY WF11 0BU
 Tel: 01132 322400
 Fax:

Fertiliser Recommendations

The phosphate and potash recommendations shown below, are those required to replace the offtake and maintain target soil indices. The larger recommended applications for soils below target index will allow the soil to build up to this target index over a number of years. Not applying fertiliser to soils which are above target index will allow the soil to run down over a number of years to the target index.

The recommendation should be increased or decreased where yields are substantially more or less than that specified. The amount to apply can be calculated using the expected yield and values for the offtake of phosphate and potash per tonne of yield given in the RB209 9th edition.

All recommendations are given for the mid-point of each Index.

Where a soil analysis value (as given by the laboratory) is close to the range of an adjacent Index, the recommendation may be reduced or increased slightly taking account of the recommendation given for the adjacent Index. Small adjustments of less than 10 kg/ha are generally not justified.

Efficient use of P and K is most likely to be achieved on soils that are well structured and enable good rooting.

For visual evaluation of soil structure (VESS), a score on 1 or 2 would be considered adequate.

Don't forget to deduct nutrients applied as organic manures.

For Nitrogen recommendations please refer to the RB209 9th edition or seek advice from an FACTS qualified adviser.

Target Indices:

Arable, Forage, Grassland and Potato Crops: P Index 2, K Index 2-

(In rotations where most crops are Autumn-sown, soils are in good condition and P is applied annually, high index 1 can be an adequate target.)

Vegetables and Bulbs: P Index 3, K Index 2+

(If vegetables are only grown occasionally as part of an arable rotation, it would be most economic to target index 2 for arable and forage crops.)

Fruit Vines and Hops: P Index 2, K Index 2, Mg Index 2

(Note: Cider apples respond to K Index 3, Mg Index 3)

A lime recommendation is usually for a 20cm depth of cultivated soil or a 15cm depth of grassland soil. Where soil is acid below 20 cm and soils are ploughed for arable crops, a proportionately larger quantity of lime should be applied. However, if more than 10 t/ha is needed, half should be deeply cultivated into the soil and ploughed down, with the remainder applied to the surface and worked in.

For established grassland or other situations where there is no, or only minimal soil cultivation, no more than 7.5 t/ha of lime should be applied in one application.

In these situations, applications of lime change the pH below the surface very slowly. Consequently, the underlying soil should not be allowed to become too acidic because this will affect the root growth and thus limit nutrient and water uptake, which will adversely affect yield.

Field Name / Ref / Soil Type	Last Crop / Next Crop	P2O5	K2O	MgO	Lime (Arable) (Grass)	
		Units/Acre			T/Ac	
		Kg/Ha			Te/Ha	
BLAENYCOED 1 025291 /	Not Given / Not Given				0	0
BLAENYCOED 2 025292 /	Not Given / Not Given				2.3	0.6
BLAENYCOED 3 025293 /	Not Given / Not Given				2.0	0
BLAENYCOED 4 025294 /	Not Given / Not Given				2.0	0
BLAENYCOED 7 025295 /	Not Given / Not Given				3.1	1.3
					7.7	3.1

Fertiliser recommendations are based on **AHDB RB209 (Ninth Edition)**. If a nutrient is deficient and no recommendation is given, either no recommendation is given in RB209 or we have insufficient data to give a recommendation. Apply Lime to the nearest half Ton / Tonne. NRM is a UKAS accredited laboratory to ISO/IEC 17025

Report continued.....



DATE 22nd April 2025
SAMPLES FROM

SAMPLED BY

Report reference 05659/25

JOLYON PRIDDING
4RECYCLING LTD
CONTROL HOUSE
A1 BUSINESS PARK
KNOTTINGLEY ROAD
KNOTTINGLEY WF11 0BU
Tel: 01132 322400
Fax:

Fertiliser Recommendations

<i>Field Name / Ref / Soil Type</i>	<i>Last Crop / Next Crop</i>	<i>P2O5</i>	<i>K2O</i>	<i>MgO</i>	<i>Lime (Arable) (Grass)</i>		
BLAENYCOED 8	Not Given / Not Given				<i>T/Ac</i>	2.3	0.6
025296 /		<i>Units/Acre</i>			<i>Te/Ha</i>	5.6	1.6
		<i>Kg/Ha</i>					

Fertiliser recommendations are based on **AHDB RB209 (Ninth Edition)**. If a nutrient is deficient and no recommendation is given, either no recommendation is given in RB209 or we have insufficient data to give a recommendation. Apply Lime to the nearest half Ton / Tonne.
NRM is a UKAS accredited laboratory to ISO/IEC 17025

Contact : JOLYON PRIDDING
 4RECYCLING LTD
 CONTROL HOUSE
 A1 BUSINESS PARK
 KNOTTINGLEY ROAD
 KNOTTINGLEY WF11 0BU
 Tel. : 01132 322400

V537

Please quote the above code for all enquiries

Client :

Sample Matrix : Agricultural Soil

Laboratory Reference

Card Number 05658/25

Date Received 14-Apr-25

Date Reported 22-Apr-25

SOIL ANALYSIS REPORT

Laboratory Sample Reference	Field Details			Index			mg/l (Available)		
	No.	Name or O.S. Reference with Cropping Details	Soil pH	P	K	Mg	P	K	Mg
25288/25	2	PENNANT 20 <i>No cropping details given</i>	6.1	3	2-	2	31.8	141	90
25289/25	3	PENNANT 21 <i>No cropping details given</i>	6.4	3	1	2	32.2	107	79
25290/25	4	PENNANT 22 <i>No cropping details given</i>	6.0	3	1	2	28.0	111	96

If general fertiliser and lime recommendations have been requested, these are given on the following sheets.

The analytical methods used are as described in DEFRA Reference Book 427

The index values are determined from the AHDB Fertiliser Recommendations RB209 9th Edition.

Released by*Katie Dunn*..... On behalf of NRM Date *22/04/25*.....

Independently Analysed by **NRM**, part of the **Cawood Group**, Coopers Bridge, Braziers Lane, Bracknell, Berkshire RG42 6NS

Tel +44 (0) 1344 886338 Fax: +44 (0) 1344 890972 Email: enquiries@nrm.uk.com www.nrm.uk.com



DATE 22nd April 2025
 SAMPLES FROM

SAMPLED BY

Report reference 05658/25

JOLYON PRIDDING
 4RECYCLING LTD
 CONTROL HOUSE
 A1 BUSINESS PARK
 KNOTTINGLEY ROAD
 KNOTTINGLEY WF11 0BU
 Tel: 01132 322400
 Fax:

Fertiliser Recommendations

The phosphate and potash recommendations shown below, are those required to replace the offtake and maintain target soil indices. The larger recommended applications for soils below target index will allow the soil to build up to this target index over a number of years. Not applying fertiliser to soils which are above target index will allow the soil to run down over a number of years to the target index.

The recommendation should be increased or decreased where yields are substantially more or less than that specified. The amount to apply can be calculated using the expected yield and values for the offtake of phosphate and potash per tonne of yield given in the RB209 9th edition.

All recommendations are given for the mid-point of each Index.

Where a soil analysis value (as given by the laboratory) is close to the range of an adjacent Index, the recommendation may be reduced or increased slightly taking account of the recommendation given for the adjacent Index. Small adjustments of less than 10 kg/ha are generally not justified.

Efficient use of P and K is most likely to be achieved on soils that are well structured and enable good rooting.

For visual evaluation of soil structure (VESS), a score on 1 or 2 would be considered adequate.

Don't forget to deduct nutrients applied as organic manures.

For Nitrogen recommendations please refer to the RB209 9th edition or seek advice from an FACTS qualified adviser.

Target Indices:

Arable, Forage, Grassland and Potato Crops: P Index 2, K Index 2-

(In rotations where most crops are Autumn-sown, soils are in good condition and P is applied annually, high index 1 can be an adequate target.)

Vegetables and Bulbs: P Index 3, K Index 2+

(If vegetables are only grown occasionally as part of an arable rotation, it would be most economic to target index 2 for arable and forage crops.)

Fruit Vines and Hops: P Index 2, K Index 2, Mg Index 2

(Note: Cider apples respond to K Index 3, Mg Index 3)

A lime recommendation is usually for a 20cm depth of cultivated soil or a 15cm depth of grassland soil. Where soil is acid below 20 cm and soils are ploughed for arable crops, a proportionately larger quantity of lime should be applied. However, if more than 10 t/ha is needed, half should be deeply cultivated into the soil and ploughed down, with the remainder applied to the surface and worked in.

For established grassland or other situations where there is no, or only minimal soil cultivation, no more than 7.5 t/ha of lime should be applied in one application.

In these situations, applications of lime change the pH below the surface very slowly. Consequently, the underlying soil should not be allowed to become too acidic because this will affect the root growth and thus limit nutrient and water uptake, which will adversely affect yield.

Field Name / Ref / Soil Type	Last Crop / Next Crop	P2O5	K2O	MgO	Lime (Arable) (Grass)
PENNANT 20	Not Given / Not Given	Units/Acre			T/Ac 1.7 0
025288 /		Kg/Ha			Te/Ha 4.2 0

Field Name / Ref / Soil Type	Last Crop / Next Crop	P2O5	K2O	MgO	Lime (Arable) (Grass)
PENNANT 21	Not Given / Not Given	Units/Acre			T/Ac 0.8 0
025289 /		Kg/Ha			Te/Ha 2.1 0

Field Name / Ref / Soil Type	Last Crop / Next Crop	P2O5	K2O	MgO	Lime (Arable) (Grass)
PENNANT 22	Not Given / Not Given	Units/Acre			T/Ac 2.0 0
025290 /		Kg/Ha			Te/Ha 4.9 0

Fertiliser recommendations are based on **AHDB RB209 (Ninth Edition)**. If a nutrient is deficient and no recommendation is given, either no recommendation is given in RB209 or we have insufficient data to give a recommendation. Apply Lime to the nearest half Ton / Tonne.
 NRM is a UKAS accredited laboratory to ISO/IEC 17025