

Contact : JOLYON PRIDDING
4RECYCLING LTD
CONTROL HOUSE
A1 BUSINESS PARK
KNOTTINGLEY ROAD
KNOTTINGLEY WF11 0BU
Tel. : 01132 322400

V537

Please quote the above code for all enquiries

Client : THE DDERW

Sample Matrix : Agricultural Soil

Laboratory Reference

Card Number 14171/24

Date Received 14-Nov-24

Date Reported 18-Nov-24

SOIL ANALYSIS REPORT

Laboratory Sample Reference	Field Details		Soil pH	Index			mg/l (Available)		
	No.	Name or O.S. Reference with Cropping Details		P	K	Mg	P	K	Mg
60654/24	1	THE DDERW 1 <i>No cropping details given</i>	5.9	1	1	2	14.6	69	69
60655/24	2	THE DDERW 2 <i>No cropping details given</i>	6.1	1	0	2	10.2	48	57
60656/24	3	THE DDERW 3 <i>No cropping details given</i>	6.6	1	2-	2	12.0	145	62
60657/24	4	THE DDERW 4 <i>No cropping details given</i>	6.3	2	1	1	16.2	64	39
60658/24	5	THE DDERW 5 <i>No cropping details given</i>	6.6	2	3	1	24.0	248	41
60659/24	6	THE DDERW 6 <i>No cropping details given</i>	6.5	3	3	2	34.2	251	51

If general fertiliser and lime recommendations have been requested, these are given on the following sheets.

The analytical methods used are as described in DEFRA Reference Book 427

The index values are determined from the AHDB Fertiliser Recommendations RB209 9th Edition.

Released by Sandy Cameron On behalf of NRM Date 18/11/24

Independently Analysed by **NRM**, part of the **Cawood Group**, Coopers Bridge, Braziers Lane, Bracknell, Berkshire RG42 6NS

Tel +44 (0) 1344 886338 Fax: +44 (0) 1344 890972 Email: enquiries@nrm.uk.com www.nrm.uk.com

DATE 18th November 2024
 SAMPLES FROM THE DDERW

JOLYON PRIDDING
 4RECYCLING LTD
 CONTROL HOUSE
 A1 BUSINESS PARK
 KNOTTINGLEY ROAD
 KNOTTINGLEY WF11 0BU
 Tel: 01132 322400
 Fax:

SAMPLED BY

Report reference 14171/24

Fertiliser Recommendations

The phosphate and potash recommendations shown below, are those required to replace the offtake and maintain target soil indices. The larger recommended applications for soils below target index will allow the soil to build up to this target index over a number of years. Not applying fertiliser to soils which are above target index will allow the soil to run down over a number of years to the target index.

The recommendation should be increased or decreased where yields are substantially more or less than that specified. The amount to apply can be calculated using the expected yield and values for the offtake of phosphate and potash per tonne of yield given in the RB209 9th edition.

All recommendations are given for the mid-point of each Index.

Where a soil analysis value (as given by the laboratory) is close to the range of an adjacent Index, the recommendation may be reduced or increased slightly taking account of the recommendation given for the adjacent Index. Small adjustments of less than 10 kg/ha are generally not justified.

Efficient use of P and K is most likely to be achieved on soils that are well structured and enable good rooting.

For visual evaluation of soil structure (VESS), a score on 1 or 2 would be considered adequate.

Don't forget to deduct nutrients applied as organic manures.

For Nitrogen recommendations please refer to the RB209 9th edition or seek advice from an FACTS qualified adviser.

Target Indices:

Arable, Forage, Grassland and Potato Crops: P Index 2, K Index 2-

(In rotations where most crops are Autumn-sown, soils are in good condition and P is applied annually, high index 1 can be an adequate target.)

Vegetables and Bulbs: P Index 3, K Index 2+

(If vegetables are only grown occasionally as part of an arable rotation, it would be most economic to target index 2 for arable and forage crops.)

Fruit Vines and Hops: P Index 2, K Index 2, Mg Index 2

(Note: Cider apples respond to K Index 3, Mg Index 3)

A lime recommendation is usually for a 20cm depth of cultivated soil or a 15cm depth of grassland soil. Where soil is acid below 20 cm and soils are ploughed for arable crops, a proportionately larger quantity of lime should be applied. However, if more than 10 t/ha is needed, half should be deeply cultivated into the soil and ploughed down, with the remainder applied to the surface and worked in.

For established grassland or other situations where there is no, or only minimal soil cultivation, no more than 7.5 t/ha of lime should be applied in one application.

In these situations, applications of lime change the pH below the surface very slowly. Consequently, the underlying soil should not be allowed to become too acidic because this will affect the root growth and thus limit nutrient and water uptake, which will adversely affect yield.

Field Name / Ref / Soil Type	Last Crop / Next Crop	P2O5	K2O	MgO	Lime (Arable)	(Grass)
THE DDERW 1 060654 /	Not Given / Not Given	Units/Acre			T/Ac 2.3	0.6
		Kg/Ha			Te/Ha 5.6	1.6
THE DDERW 2 060655 /	Not Given / Not Given	Units/Acre			T/Ac 1.7	0
		Kg/Ha			Te/Ha 4.2	0
THE DDERW 3 060656 /	Not Given / Not Given	Units/Acre			T/Ac 0	0
		Kg/Ha			Te/Ha 0	0
THE DDERW 4 060657 /	Not Given / Not Given	Units/Acre			T/Ac 1.1	0
		Kg/Ha			Te/Ha 2.8	0
THE DDERW 5 060658 /	Not Given / Not Given	Units/Acre			T/Ac 0	0
		Kg/Ha			Te/Ha 0	0

Fertiliser recommendations are based on AHDB RB209 (Ninth Edition). If a nutrient is deficient and no recommendation is given, either no recommendation is given in RB209 or we have insufficient data to give a recommendation. Apply Lime to the nearest half Ton / Tonne. NRM is a UKAS accredited laboratory to ISO/IEC 17025

Report continued.....



DATE 18th November 2024
SAMPLES FROM THE DDERW

SAMPLED BY

Report reference 14171/24

JOLYON PRIDDING
4RECYCLING LTD
CONTROL HOUSE
A1 BUSINESS PARK
KNOTTINGLEY ROAD
KNOTTINGLEY WF11 0BU
Tel: 01132 322400
Fax:

Fertiliser Recommendations

<i>Field Name / Ref / Soil Type</i>	<i>Last Crop / Next Crop</i>	<i>P2O5</i>	<i>K2O</i>	<i>MgO</i>	<i>Lime (Arable) (Grass)</i>		
THE DDERW 6	Not Given / Not Given	<i>Units/Acre</i>			<i>T/Ac</i>	0	0
060659 /		<i>Kg/Ha</i>			<i>Te/Ha</i>	0	0

Fertiliser recommendations are based on **AHDB RB209 (Ninth Edition)**. If a nutrient is deficient and no recommendation is given, either no recommendation is given in RB209 or we have insufficient data to give a recommendation. Apply Lime to the nearest half Ton / Tonne.
NRM is a UKAS accredited laboratory to ISO/IEC 17025

Contact : JOLYON PRIDDING
4RECYCLING LTD
CONTROL HOUSE
A1 BUSINESS PARK
KNOTTINGLEY ROAD
KNOTTINGLEY WF11 0BU
Tel. : 01132 322400

V537

Please quote the above code for all enquiries

Client : THE DDERW

Sample Matrix : Agricultural Soil

Laboratory Reference

Card Number 14172/24

Date Received 14-Nov-24

Date Reported 18-Nov-24

SOIL ANALYSIS REPORT

Laboratory Sample Reference	Field Details			Index			mg/l (Available)		
	No.	Name or O.S. Reference with Cropping Details	Soil pH	P	K	Mg	P	K	Mg
60660/24	1	THE DDERW 7 <i>No cropping details given</i>	5.9	4	2+	2	58.8	218	58
60661/24	2	THE DDERW 8 <i>No cropping details given</i>	5.8	2	2-	3	24.6	170	102

If general fertiliser and lime recommendations have been requested, these are given on the following sheets.

The analytical methods used are as described in DEFRA Reference Book 427

The index values are determined from the AHDB Fertiliser Recommendations RB209 9th Edition.

Released by *Sandy Cameron* On behalf of NRM Date *18/11/24*

Independently Analysed by **NRM**, part of the **Cawood Group**, Coopers Bridge, Braziers Lane, Bracknell, Berkshire RG42 6NS

Tel +44 (0) 1344 886338 Fax: +44 (0) 1344 890972 Email: enquiries@nrm.uk.com www.nrm.uk.com



DATE 18th November 2024
SAMPLES FROM THE DDERW

SAMPLED BY

Report reference 14172/24

JOLYON PRIDDING
4RECYCLING LTD
CONTROL HOUSE
A1 BUSINESS PARK
KNOTTINGLEY ROAD
KNOTTINGLEY WF11 0BU
Tel: 01132 322400
Fax:

Fertiliser Recommendations

The phosphate and potash recommendations shown below, are those required to replace the offtake and maintain target soil indices. The larger recommended applications for soils below target index will allow the soil to build up to this target index over a number of years. Not applying fertiliser to soils which are above target index will allow the soil to run down over a number of years to the target index.

The recommendation should be increased or decreased where yields are substantially more or less than that specified. The amount to apply can be calculated using the expected yield and values for the offtake of phosphate and potash per tonne of yield given in the RB209 9th edition.

All recommendations are given for the mid-point of each Index.

Where a soil analysis value (as given by the laboratory) is close to the range of an adjacent Index, the recommendation may be reduced or increased slightly taking account of the recommendation given for the adjacent Index. Small adjustments of less than 10 kg/ha are generally not justified.

Efficient use of P and K is most likely to be achieved on soils that are well structured and enable good rooting.

For visual evaluation of soil structure (VESS), a score on 1 or 2 would be considered adequate.

Don't forget to deduct nutrients applied as organic manures.

For Nitrogen recommendations please refer to the RB209 9th edition or seek advice from an FACTS qualified adviser.

Target Indices:

Arable, Forage, Grassland and Potato Crops: P Index 2, K Index 2-

(In rotations where most crops are Autumn-sown, soils are in good condition and P is applied annually, high index 1 can be an adequate target.)

Vegetables and Bulbs: P Index 3, K Index 2+

(If vegetables are only grown occasionally as part of an arable rotation, it would be most economic to target index 2 for arable and forage crops.)

Fruit Vines and Hops: P Index 2, K Index 2, Mg Index 2

(Note: Cider apples respond to K Index 3, Mg Index 3)

A lime recommendation is usually for a 20cm depth of cultivated soil or a 15cm depth of grassland soil. Where soil is acid below 20 cm and soils are ploughed for arable crops, a proportionately larger quantity of lime should be applied. However, if more than 10 t/ha is needed, half should be deeply cultivated into the soil and ploughed down, with the remainder applied to the surface and worked in.

For established grassland or other situations where there is no, or only minimal soil cultivation, no more than 7.5 t/ha of lime should be applied in one application.

In these situations, applications of lime change the pH below the surface very slowly. Consequently, the underlying soil should not be allowed to become too acidic because this will affect the root growth and thus limit nutrient and water uptake, which will adversely affect yield.

Field Name / Ref / Soil Type	Last Crop / Next Crop	P2O5	K2O	MgO	Lime (Arable)	(Grass)
THE DDERW 7	Not Given / Not Given	Units/Acre			T/Ac 2.3	0.6
060660 /		Kg/Ha			Te/Ha 5.6	1.6

Field Name / Ref / Soil Type	Last Crop / Next Crop	P2O5	K2O	MgO	Lime (Arable)	(Grass)
THE DDERW 8	Not Given / Not Given	Units/Acre			T/Ac 2.5	0.8
060661 /		Kg/Ha			Te/Ha 6.3	2.1

Fertiliser recommendations are based on **AHDB RB209 (Ninth Edition)**. If a nutrient is deficient and no recommendation is given, either no recommendation is given in RB209 or we have insufficient data to give a recommendation. Apply Lime to the nearest half Ton / Tonne.

NRM is a UKAS accredited laboratory to ISO/IEC 17025