

Contact : MATTHEW CONOLLY
 4R GROUP
 12C NEWENT BUS PARK
 GLOUCESTER STREET
 NEWENT
 GLOUCESTERSHIRE
 GL18 1DZ
 Tel. : 07508322259

V293

Please quote the above code for all enquiries

Client : GARETH PARRY
 BRYN DROFA
 LLANBABO
 RHOSGOCH
 ANGLESEY
 LL66 0BN

Sample Matrix : Agricultural Soil

Laboratory Reference

Card Number 03873/26

Date Received 13-Apr-26

Date Reported 23-Apr-26

SOIL ANALYSIS REPORT

Laboratory Sample Reference	Field Details			Index			mg/l (Available)		
	No.	Name or O.S. Reference with Cropping Details	Soil pH	P	K	Mg	P	K	Mg
16435/26	1	FIELD 1 <i>No cropping details given</i>	8.2	3	2-	2	33.0	157	88
16436/26	2	FIELD 2 <i>No cropping details given</i>	8.1	3	2+	2	35.4	206	90
16437/26	3	FIELD 3 <i>No cropping details given</i>	8.4	2	2-	2	24.6	145	97
16438/26	4	FIELD 4 <i>No cropping details given</i>	8.4	2	2-	2	24.8	147	83

If general fertiliser and lime recommendations have been requested, these are given on the following sheets.

The analytical methods used are as described in DEFRA Reference Book 427

The index values are determined from the AHDB Fertiliser Recommendations RB209 9th Edition.

Released by*Katie Dunn*..... On behalf of NRM Date*23/04/26*.....

Independently Analysed by **NRM**, part of the **Cawood Group**, Coopers Bridge, Braziers Lane, Bracknell, Berkshire RG42 6NS

Tel +44 (0) 1344 886338 Fax: +44 (0) 1344 890972 Email: enquiries@nrm.uk.com www.nrm.uk.com



DATE 23rd April 2026
 SAMPLES FROM GARETH PARRY, BRYN DROFA,
 LLANBABO, RHOSGOCH

MATTHEW CONOLLY
 4R GROUP
 12C NEWENT BUS PARK
 GLOUCESTER STREET
 NEWENT
 GLOUCESTERSHIRE
 Tel: 0750832259
 Fax:

SAMPLED BY

Report reference 03873/26

Fertiliser Recommendations

The phosphate and potash recommendations shown below, are those required to replace the offtake and maintain target soil indices. The larger recommended applications for soils below target index will allow the soil to build up to this target index over a number of years. Not applying fertiliser to soils which are above target index will allow the soil to run down over a number of years to the target index.

The recommendation should be increased or decreased where yields are substantially more or less than that specified. The amount to apply can be calculated using the expected yield and values for the offtake of phosphate and potash per tonne of yield given in the RB209 9th edition.

All recommendations are given for the mid-point of each Index.

Where a soil analysis value (as given by the laboratory) is close to the range of an adjacent Index, the recommendation may be reduced or increased slightly taking account of the recommendation given for the adjacent Index. Small adjustments of less than 10 kg/ha are generally not justified.

Efficient use of P and K is most likely to be achieved on soils that are well structured and enable good rooting.

For visual evaluation of soil structure (VESS), a score on 1 or 2 would be considered adequate.

Don't forget to deduct nutrients applied as organic manures.

For Nitrogen recommendations please refer to the RB209 9th edition or seek advice from an FACTS qualified adviser.

Target Indices:

Arable, Forage, Grassland and Potato Crops: P Index 2, K Index 2-

(In rotations where most crops are Autumn-sown, soils are in good condition and P is applied annually, high index 1 can be an adequate target.)

Vegetables and Bulbs: P Index 3, K Index 2+

(If vegetables are only grown occasionally as part of an arable rotation, it would be most economic to target index 2 for arable and forage crops.)

Fruit Vines and Hops: P Index 2, K Index 2, Mg Index 2

(Note: Cider apples respond to K Index 3, Mg Index 3)

A lime recommendation is usually for a 20cm depth of cultivated soil or a 15cm depth of grassland soil. Where soil is acid below 20 cm and soils are ploughed for arable crops, a proportionately larger quantity of lime should be applied. However, if more than 10 t/ha is needed, half should be deeply cultivated into the soil and ploughed down, with the remainder applied to the surface and worked in.

For established grassland or other situations where there is no, or only minimal soil cultivation, no more than 7.5 t/ha of lime should be applied in one application.

In these situations, applications of lime change the pH below the surface very slowly. Consequently, the underlying soil should not be allowed to become too acidic because this will affect the root growth and thus limit nutrient and water uptake, which will adversely affect yield.

Field Name / Ref / Soil Type	Last Crop / Next Crop	P2O5	K2O	MgO	Lime (Arable)	(Grass)
FIELD 1	Not Given / Not Given	Units/Acre			T/Ac	0
016435 /		Kg/Ha			Te/Ha	0
FIELD 2	Not Given / Not Given	Units/Acre			T/Ac	0
016436 /		Kg/Ha			Te/Ha	0
FIELD 3	Not Given / Not Given	Units/Acre			T/Ac	0
016437 /		Kg/Ha			Te/Ha	0
FIELD 4	Not Given / Not Given	Units/Acre			T/Ac	0
016438 /		Kg/Ha			Te/Ha	0

Fertiliser recommendations are based on **AHDB RB209 (Ninth Edition)**. If a nutrient is deficient and no recommendation is given, either no recommendation is given in RB209 or we have insufficient data to give a recommendation. Apply Lime to the nearest half Ton / Tonne.

NRM is a UKAS accredited laboratory to ISO/IEC 17025

Contact : MATTHEW CONOLLY
 4R GROUP
 12C NEWENT BUS PARK
 GLOUCESTER STREET
 NEWENT
 GLOUCESTERSHIRE
 GL18 1DZ
 Tel. : 07508322259

V293

Please quote the above code for all enquiries

Client : ARWEL HUGHES
 PLAS NEWYDD
 LLANBABO
 RHOSGOCH
 ANGLESEY
 LL66 0BN

Sample Matrix : Agricultural Soil

Laboratory Reference

Card Number 03872/26

Date Received 13-Apr-26

Date Reported 23-Apr-26

SOIL ANALYSIS REPORT

Laboratory Sample Reference	Field Details			Index			mg/l (Available)		
	No.	Name or O.S. Reference with Cropping Details	Soil pH	P	K	Mg	P	K	Mg
16433/26	1	FIELD 1 <i>No cropping details given</i>	8.1	3	2-	2	27.0	155	90
16434/26	2	FIELD 2 <i>No cropping details given</i>	8.2	3	2-	2	34.8	159	84

If general fertiliser and lime recommendations have been requested, these are given on the following sheets.

The analytical methods used are as described in DEFRA Reference Book 427

The index values are determined from the AHDB Fertiliser Recommendations RB209 9th Edition.

Released by*Katie Dunn*..... On behalf of NRM Date*23/04/26*.....

Independently Analysed by **NRM**, part of the **Cawood Group**, Coopers Bridge, Braziers Lane, Bracknell, Berkshire RG42 6NS

Tel +44 (0) 1344 886338 Fax: +44 (0) 1344 890972 Email: enquiries@nrm.uk.com www.nrm.uk.com



DATE 23rd April 2026
 SAMPLES FROM ARWEL HUGHES, PLAS NEWYDD,
 LLANBABO, RHOSGOCH

MATTHEW CONOLLY
 4R GROUP
 12C NEWENT BUS PARK
 GLOUCESTER STREET
 NEWENT
 GLOUCESTERSHIRE
 Tel: 07508322259
 Fax:

SAMPLED BY

Report reference 03872/26

Fertiliser Recommendations

The phosphate and potash recommendations shown below, are those required to replace the offtake and maintain target soil indices. The larger recommended applications for soils below target index will allow the soil to build up to this target index over a number of years. Not applying fertiliser to soils which are above target index will allow the soil to run down over a number of years to the target index.

The recommendation should be increased or decreased where yields are substantially more or less than that specified. The amount to apply can be calculated using the expected yield and values for the offtake of phosphate and potash per tonne of yield given in the RB209 9th edition.

All recommendations are given for the mid-point of each Index.

Where a soil analysis value (as given by the laboratory) is close to the range of an adjacent Index, the recommendation may be reduced or increased slightly taking account of the recommendation given for the adjacent Index. Small adjustments of less than 10 kg/ha are generally not justified.

Efficient use of P and K is most likely to be achieved on soils that are well structured and enable good rooting.

For visual evaluation of soil structure (VESS), a score on 1 or 2 would be considered adequate.

Don't forget to deduct nutrients applied as organic manures.

For Nitrogen recommendations please refer to the RB209 9th edition or seek advice from an FACTS qualified adviser.

Target Indices:

Arable, Forage, Grassland and Potato Crops: P Index 2, K Index 2-

(In rotations where most crops are Autumn-sown, soils are in good condition and P is applied annually, high index 1 can be an adequate target.)

Vegetables and Bulbs: P Index 3, K Index 2+

(If vegetables are only grown occasionally as part of an arable rotation, it would be most economic to target index 2 for arable and forage crops.)

Fruit Vines and Hops: P Index 2, K Index 2, Mg Index 2

(Note: Cider apples respond to K Index 3, Mg Index 3)

A lime recommendation is usually for a 20cm depth of cultivated soil or a 15cm depth of grassland soil. Where soil is acid below 20 cm and soils are ploughed for arable crops, a proportionately larger quantity of lime should be applied. However, if more than 10 t/ha is needed, half should be deeply cultivated into the soil and ploughed down, with the remainder applied to the surface and worked in.

For established grassland or other situations where there is no, or only minimal soil cultivation, no more than 7.5 t/ha of lime should be applied in one application.

In these situations, applications of lime change the pH below the surface very slowly. Consequently, the underlying soil should not be allowed to become too acidic because this will affect the root growth and thus limit nutrient and water uptake, which will adversely affect yield.

Field Name / Ref / Soil Type	Last Crop / Next Crop	P2O5	K2O	MgO	Lime (Arable) (Grass)		
FIELD 1	Not Given / Not Given	Units/Acre			T/Ac	0	0
016433 /		Kg/Ha			Te/Ha	0	0
FIELD 2	Not Given / Not Given	Units/Acre			T/Ac	0	0
016434 /		Kg/Ha			Te/Ha	0	0

Fertiliser recommendations are based on **AHDB RB209 (Ninth Edition)**. If a nutrient is deficient and no recommendation is given, either no recommendation is given in RB209 or we have insufficient data to give a recommendation. Apply Lime to the nearest half Ton / Tonne.

NRM is a UKAS accredited laboratory to ISO/IEC 17025